

ME 2580 Example #3: (Rectilinear Motion)

Given: $a(v) = g - cv$ ($a(v) \geq 0$) ... the **acceleration** of an object falling in a fluid.
initial conditions: $v(0) = 0$

Find: $v(t)$... the **velocity** of the particle as a **function of time**

Solution:

$$a(v) = \frac{dv}{dt} = g - cv \Rightarrow \frac{dv}{g - cv} = dt \Rightarrow \frac{-1}{c} \int \frac{-c dv}{g - cv} = \int dt$$

using indefinite integrals

$$\frac{-1}{c} \ln(g - cv) = t + D$$

Aside:

$$\int \frac{f'(x) dx}{f(x)} = \ln(f(x))$$

Applying the **initial condition**, $v(0) = 0$, gives $D = \frac{-1}{c} \ln(g)$

Substituting the value of the constant D into the first boxed equation and **simplifying**:

$$\frac{-1}{c} \ln(g - cv) = t + \frac{-1}{c} \ln(g) \Rightarrow \ln(g - cv) = \ln(g) - ct \Rightarrow g - cv = \exp(\ln(g) - ct)$$

$$g - cv = \exp(\ln(g)) \cdot \exp(-ct) = g e^{-ct} \Rightarrow v(t) = \frac{g}{c} (1 - e^{-ct})$$

The function $v(t)$ starts at zero and increases exponentially to the **final value** of $\frac{g}{c}$. The **larger** the value of the coefficient c , the **faster** the function increases toward $\frac{g}{c}$. The plot below shows a plot a $v(t)$ for $c = g = 1$.

